

Original Research

Utilization of Pomegranate Peel Pectin as a Functional Ingredient for Nutritive Jelly Formulation: A Step Toward Healthier Processed Food

Trishna Roy¹, Nilufa Yeasmin^{1,*}, Ayesha Begum^{1,*}, Md. Altaf Hossain^{1,2}, Mohammad Mozibul Haque¹, Kanij Fatema Nishan¹, Md. Zia Uddin Al Mamun³, Anjum Mahfuza⁴ and Sultana Jannat Pomy¹

¹Department of Applied Food Science and Nutrition, Faculty of Food Science and Technology, Chattogram Veterinary and Animal Sciences University (CVASU), Khulshi, Chattogram-4225, Bangladesh

²Department of Food Science, University of Otago, Dunedin 9054, New Zealand

³Institute of Food Science and Technology (IFST), Bangladesh Council of Scientific and Industrial Research (BCSIR), Dhaka-1205, Bangladesh

⁴Department of Environmental Science, Kunming University of Science and Technology, Kunming, Yunnan, China

Article history

Received: 1 August 2025

Revised: 19 September 2025

Accepted: 23 September 2025

Published Online: 30 September 2025

*Correspondence:

Ayesha Begum; Nilufa Yeasmin

Address: Department of Applied Food Science and Nutrition, Faculty of Food Science and Technology, Chattogram Veterinary and Animal Sciences University, Khulshi, Chattogram-4225, Bangladesh.

Email: asha03du@gmail.com (AB);
lily.fst2010@gmail.com (NY)

How to cite this article: Roy T, Yeasmin N, Begum A, Hossain MA, Haque MM, Nishan KF, Al Mamun MZU, Mahfuza A, Pomy SJ. Utilization of Pomegranate Peel Pectin as a Functional Ingredient for Nutritive Jelly Formulation: A Step Toward Healthier Processed Food. *Health Dynamics*, 2025, 2(9), 368-379.
<https://doi.org/10.33846/hd20901>



Copyrights: © 2025 by the authors. This is an open access article under the terms and conditions of the Creative Commons Attribution – NoDerivatives 4.0 International (CC BY-ND 4.0) license (<https://creativecommons.org/licenses/by-nd/4.0/>).

ABSTRACT

Background: The growing prevalence of chronic diseases has increased interest in natural functional foods over synthetic alternative. Pomegranate peel pectin, a natural gelling agent with dietary fiber, bioactive substances that helps with better digestion, glycemic control and antioxidant defense. As a healthy substitute for commercial pectin, the current study aimed to extract pectin from pomegranate peel and assess its stability and functionality in jelly preparation. **Methods:** Pectin was extracted from pomegranate peel powder using citric acid. Fourier Transform Infrared (FT-IR) spectroscopy was used to investigate structural characteristics. Galacturonic acid, amidation, ash content, degree of methyl esterification, and water holding capacity of the extracted pectin were all measured. Pomegranate juice's proximate composition was ascertained. Both commercial and extracted pectin were used to make the jellies, sweetened with sugar or honey and tested for proximate composition, fiber content, antioxidant activity, sensory quality, and microbiological safety while being stored. **Results:** The extracted pectin yield was 8.2%, with a high methoxyl content, 1.15% ash, and 235.25% water-holding capacity. Pomegranate juice contained 85.3% moisture, 10.5%, total sugars, 0.15g citric acid, 0.9g ascorbic acid, and 0.03g ash. Jellies made with extracted pectin showed similar sensory and proximate property to those made with commercial pectin. Fiber content increased in jellies with extracted pectin, particularly with honey. Microbial investigation found sugar-added jelly safe for two months and honey-added for three months when refrigerated. **Conclusion:** This study shows pomegranate peel pectin, a health promoting organic substitute for commercial pectin, can be effectively used in jelly formulation.

Keywords: Pomegranate peel pectin; functional food; jelly formulation; antioxidant activity; dietary fiber; food preservation

1. INTRODUCTION

In recent years, the food market has witnessed a consistent emergence of new trend due to consumers' evolving tastes for healthier food options. The growing interest in natural alternative that ensures both products quality and functional health benefits rather than processed food has led to a focus on the use of functional ingredients that are safe and offer potential medicinal and nutritional benefits.^(1,2)

Punica granatum L., which is part of the Punicaceae family, has been regarded as a type of functional fruit due to evidence of its health-promoting qualities from several studies, primarily ascribed to punicalagin and, to a lesser extent, other metabolites, like flavonols and anthocyanin.^(3,4) Bioactive phytochemicals found in pomegranate fruit, flowers, bark and leaves have antimicrobial properties, lower blood pressure, and prevent major diseases like diabetes and cancer. These results validate the traditional use of the pomegranates as a medicine.⁽⁵⁾ Despite its phytochemical richness, approximately 30-40% of the fruit's weight is wasted as peel during juice preparation. This by-product contains an abundance of polyphenols, flavonoids, dietary fiber, and pectin, all associated with antioxidant, anti-inflammatory, and metabolic health benefits.⁽⁶⁾

According to Abid et al. (2017), the pomegranate peel is a major non-edible part that contains about 10% pectin, making it a promising and novel pectin source.⁽⁷⁾ Pectin, a natural heteropolysaccharide utilized as a gelling agent, has drawn interest for both its technological applications and its therapeutic potential. Recent studies evidences its capacity to reduce cholesterol, regulate blood glucose, enhance gut health, and prevent cancer cell aggregation, establishing it as a highly valuable biofunctional food component.⁽⁸⁻¹⁰⁾

There have been multiple reports of functional food products containing compounds from pomegranate peel. Incorporating peel extracts into nutraceutical gummies and jellies have been found to increase antioxidant activity and polyphenol bioaccessibility,^(11,12) while using pectin derived from the peel as a fat substitute in confectionaries has improved oxidative stability.⁽¹³⁾ Pectin produced from citrus fruits has also demonstrated promising as a natural emulsifier in dairy systems and in bioactive films that can prolong the shelflife of food.^(14,15)

There being an increasing interest in the use of fruit-by products for functional food production, there is limited study that specifically investigates pectin derived from pomegranate peel as the principle gelling ingredient in jelly formulation. This gap limits the implementation of pomegranate peel pectin in practical uses that could increase product texture and provide health advantages including antioxidant activity, dietary fiber enrichment, and metabolic regulation. Bridging this knowledge gap would enhance the development of healthier confectionary products and facilitate the sustainable utilization of fruit by-products while processing. Therefore, this study aims to investigate the potential of pomegranate peel pectin as a functional food ingredient by its extraction, characterization, and application in jelly formulation.

2. METHODS

2.1 Materials and Reagents

Fresh pomegranates (*Punica Granatum* L) were obtained from local markets in Chattogram, Bangladesh. All analytical grade reagents and chemical such as hydrochloric acid (HCl), sodium hydroxide (NaOH), ethanol, acetone, citric acid and phenol-sulfuric acid, 2,2-Diphenyl-1-picryl hydrazyl (DPPH), were acquired from Sigma-Aldrich Co. (St. Louis, Missouri, USA) for the extraction and characterization of pectin.

For jelly preparation, ingredients such as sugar, honey, and Commercial grade fruit pectin were procured from local grocery stores. All materials were handled with proper hygienic practices to ensure quality and safety throughout the study.

2.2 Sample Collection and Preparation of Pomegranate Peel Powder

Mature and good-quality pomegranate fruits were purchased from Chattogram City Corporation's community market. Collected fruits were washed and peeled manually. To deactivate the pectinase enzyme, pomegranate peels were immersed in a hot water bath (90°C) for two minutes. After being dried in an air cabinet drier (Bosch MKM 600, Germany) for 48 hours at 60°C until achieving a consistent weight, the peels were ground into a powder and stored at 4°C for the next step. Figure 1 depicts the entire experimental procedure employed in this study.

2.3 Extraction of Pectin from Pomegranate Peel Powder by Citric Acid Based Aqueous Extraction

A modified version of Sathish et al. (2018) method, pectin was extracted combining 10gm of sample with 150 mL of distilled water.⁽¹⁶⁾ After that, the pH was reduced to approximately 3 using citric acid. After 25 minutes of boiling, the sample was filtered to remove peel. An equal amount of ethanol was added to the filtered extract, and it was then let to stand for an hour. The extracted pectin was separated by filtering, then precipitated with 96% ethanol and treated with 1ml of acetone. Following separation, the pectin was dried for 24 hours at 50°C before being kept at 4°C.

2.4 Physicochemical Analysis of Pectin Derived from Pomegranate Peel

To determine the effectiveness of the dried pectin extracted from pomegranate peel, a range of qualitative (Color, solubility in different solvent) and quantitative (Moisture, ash, water holding capacity, degree of esterification, amidation, galacturonic acid content, pH, and FT-IR analysis) tests were carried out.

2.4.1 Qualitative test of pomegranate peel pectin

The color was assessed visually. The solubility of dry pectin in different solvents was identified using the method provided by Aina et al. (2012).⁽¹⁷⁾




Figure 1. Schematic flow of overall experimental procedure

2.4.2 Quantitative analysis of pomegranate peel pectin

For the quantitative test, pH was measured with a pH meter (model: HI-98107, Hanna Instrument, Italy). Moisture percentage and ash content of dry pectin were determined following AOAC (2000) procedures.⁽¹⁸⁾ Using the Joint Expert Committee on Food Additives of the FAO and WHO (2009) procedure, the amidation, esterification degree, and glucuronic acid level of pectin was determined.⁽¹⁹⁾ The method provided by Zhang et al. (2017) was used to determine the liquid holding capacity (LHC) of pectin.⁽²⁰⁾

2.4.3 Fourier-Transform Infrared (FT-IR) spectroscopic analysis

FT-IR analysis of pectin by potassium bromide (KBr) pellet method was conducted using a Jasco FT/IR430 spectrophotometer (Japan) to ascertain the presence of functional group. 2mg of pectin samples were mixed with 300mg of dry KBr crystal and crushed into pellets with a rotary vacuum pump. The KBr pellet was subsequently analyzed using FT-IR spectrophotometer within 400-4000cm⁻¹ spectral range.⁽²¹⁾

2.5 Preparation of Jelly Using Pomegranate Fruit Juice and Extracted Peel Pectin

Uniformly ripe pomegranate fruits were picked, cut into two to four pieces, crushed in a grinder, and the resulting bulk was filtered through muslin cloth to collect and measure the juice for further analysis and jelly preparation. The juice's pH was measured using an HI-98107 pH meter (Hanna Instrument, Italy), and its moisture, ash, vitamin C, and titratable acidity were determined following the standard procedures of the Association of Official Analytical Chemists (2000).⁽¹⁸⁾

Jelly was prepared in four varieties using pomegranate juice, pectin extracted from pomegranate

peel, sugar, and honey in different percentages and was designated as Jelly T1, T2, T3, and T4. Table 1 displays the different treatment compositions of jelly. Pomegranate juice along with sugar or honey was simmered for 5 minutes over medium heat while being constantly stirred. Pectin and citric acid were then added to alter the pH level. While stirring, the heating was kept up. A refractometer reading of 65–67 °Brix TSS in the combination after 15 minutes indicated the endpoint. Subsequently, the jelly was transferred into a sterilized glass, appropriately labeled, and stored under freezing conditions for further experimental analysis.

Table 1. List of treatments

Treatment	Sample	Treatment combination
T ₀	PJ _a (Control)	Pomegranate jelly made with 1g commercial Pectin
T ₁	PJ _b	Sugar 40% + Pomegranate fruit juice 60% + 1g Extracted peel pectin
T ₂	PJ _c	Honey 40% + Pomegranate fruit juice 60% + 1g Extracted peel pectin
T ₃	PJ _d	Honey 50% + Pomegranate fruit juice 50% + 1g Extracted peel pectin
T ₄	PJ _e	Sugar 50% + Pomegranate fruit juice 50% + 1g Extracted peel pectin

T= Treatment; PJ= Pomegranate peel pectin jelly

2.6 Physicochemical Analysis of Jelly

The prepared jelly's physical characteristics, including pH and total soluble solids, were measured. Total soluble solids were investigated with an Atego RX 1000 digital refractometer and pH was determined with a pH meter (model: HI-98107, Hanna Instrument, Italy). The Association of Official Analytical Chemists (2000) recommended technique was used to do moisture, crude fibre, ash, fat, protein, carbohydrate analysis on the jelly.⁽¹⁸⁾

2.7 Antioxidant Activity of Jelly by DPPH Scavenging Assay

With slight modifications, Azlim et al.'s (2010) DPPH test was implemented to measure antioxidant capability.⁽²²⁾ 1gm of sample was placed in the Felcon tube for extract preparation. Following this, 10 mL of 100% methanol was added and the resultant mixture was let to sit for 72 hours. Continuous straining was done out every 4 hours. Filtration was followed by the collection of the methanolic extract. The DPPH solution was made by dissolving 6 mg of DPPH in 100 mL of methanol. For the assay, 1 mL of each extract was combined with 2 mL of the methanolic DPPH solution, and the mixture was left to incubate for 30 minutes in the dark. At 517 nm,

absorbance was taken three times with a UV-VIS spectrophotometer (UV-2600, Shimadzu Corporation, USA). Methanol by itself was utilized as the blank, while a combination of 1 mL methanol and 2 mL DPPH solution was utilized as the control. Trolox served as the standard of reference. Milligrams of Trolox equivalent per gram of extract (mg TE/g) was used to express the antioxidant capacity, which reflected the sample's power to scavenge DPPH free radicals.

The following formula was used to determine the free radical scavenging capacity of DPPH activity:

$$\% \text{ inhibition} = \frac{(\text{Abs.1} - \text{Abs.2})}{\text{Abs.1}} \times 100$$

Where, Abs.1 = Absorbance of control and Abs.2 = Absorbance of sample.

2.8 Microbiological Analysis of Jelly

Microbial quality was determined via monitoring fungal growth, fecal coliform (*Escherichia coli*). Fungal contamination was investigated using the standard plate count method on Sabouraud Dextrose Agar (SDA) and cultured at 25-30 °C for upto six weeks, where yeasts appeared creamy and mold as multicolored filaments.^(23,24) Fecal coliforms were detected using mFC agar supplemented with rosolic acid. Blue colonies after 24 hours incubation at 44.5 °C indicated presumptive *E. coli*. For confirmation, presumptive colonies were

subcultured on MacConkey agar, incubated at 37 ± 0.5 °C, and then inspected under long-wave UV light, where fluorescence indicated the presence of *E. coli*, with UV lamp accuracy verified beforehand to ensure reliable detection.⁽²⁵⁾

2.9 Sensory Evaluation

The developed pomegranate peel-pectin jelly's acceptance among consumers was evaluated by a group of ten panelists. The panelists were chosen at random from among students, faculty, and staff from the Applied Food Science and Nutrition department of Chattogram Veterinary and Animal Sciences University (CVASU). On a 1-to-9-point hedonic scale, panelists were asked to score the distinctiveness of the five formulations of jelly's appearance, color, smell, taste, sweetness, texture, and overall acceptability. The scale is set up as follows: 9 = Like Extremely, 8 = Like very much, 7 = Moderately liked, 6 = Like highly, 5 = Neither like nor dislike, 4 = Slightly dislike, 3 = Dislike moderately, 2 = Very much dislike, and 1 = Dislike extremely. The panelists were provided with water to rinse their mouths between tastings.

2.10 Statistical Analysis

Data were compiled using Microsoft Excel 2013 and analyzed using IBM SPSS Statistics version 25. Descriptive statistics were calculated for the sensory attributes and proximate composition. One-way analysis of variance (ANOVA) was performed to assess significant differences in proximate composition, phytochemical content, antioxidant capacity, and sensory evaluation among the samples. Statistical significance was determined at the 95% confidence level ($p < 0.05$). Results are expressed as mean \pm standard deviation based on three independent replicates.

2.11 Ethical Practices

This study utilized panelists from the Department of Applied Food Science and Nutrition of CVASU. Before sensory evaluation, expert assessment confirmed the jelly's safety and non-hazardous production. The sensory evaluation was performed with all participants informed about the nature of the study. Every participant was made aware of the purpose of the study before the sensory evaluation was conducted. The panel was not compensated for their voluntary participation. Participants gave their verbal consent, and their anonymity and confidentiality were strictly maintained.

3. RESULTS AND DISCUSSION

3.1 Pectin Yield of Pomegranate Peel

As a control, we utilized apple pectin powder from Triveni Interchem Pvt. Ltd. Pomegranate peel (PP) pectin yields were found to be 8.2%. The pectin yields of the peels analyzed in this study were higher than those reported in the literature, which ranged from 3.62 to 4.48%; from 11.46 to 22.09% for citrus peels (lemon, mandarin, orange and grapefruit); from 2.8% and 8.8% for banana and mango peels; from 1.99 and 2.86% for quince.⁽²⁶⁻²⁸⁾ It highlights the potential of pomegranate peel as an excellent natural source of pectin possess extra health advantages beyond its functional purpose.

3.2 Physiochemical Analysis of Pectin Extracted from Pomegranate Peel

3.2.1 Qualitative parameters of pomegranate peel pectin

The pectin extracted from pomegranate peel was grayish to light brown. Table 2 displays its solubility in different solvents. Pectin dissolved in NaOH (1M) more readily than it did in acetone and methanol. A study on lemon, grapefruit and orange peel pectin solubility resulted in the formation of yellow-colored solutions under cold alkaline conditions (20°C, 0.1 N NaOH), whereas milky solutions were formed in hot alkaline conditions (85–95 °C, 0.1 N NaOH).⁽¹⁷⁾

Table 2. Solubility of pomegranate peel pectin in different solvents

Solvent	Solubility
NaOH	Soluble
Methanol	Not soluble
Acetone	Not soluble
Hot water	Soluble
Cold water	Not soluble

3.2.2 Quantitative parameters of pomegranate peel pectin

The quantitative parameters of physiochemical analysis of pectin derived from pomegranate peel identified numerous significant factors that reflects its purity and functional characteristics. The pH of the extracted pectin was recorded at 4.2, indicating a somewhat acidic characteristic. Figure 2a delineates the moisture content, ash content, esterification degree (DE), amidation, and galacturonic acid (GA) content. The ash content measured 1.15%, signifying a minimal presence of inorganic contaminants, which is advantageous for

high quality pectin. Because, the purity of the pectin increases as the ash content decreases.⁽²⁹⁾

The degree of esterification (DE) was measured to be 57.74, classifying the isolated pectin as a high methoxyl pectin which is consistent with findings in previous studies.^(26,30) Pomegranate pectin's amidation level was 3.75%, and galacturonic acid (GA) content was 78.48%. The results showed that extracted pectin had larger quantities of GA than gold kiwifruit pectin (28.96% to 58.57%) when they were compared to those of prior studies, but pomegranate peel derived pectin had higher amidation degrees than orange and lemon peel pectin (2.20% and 1.44%, respectively).^(26,30) Pectin with high GA offers several health benefits. Galacturonic acid is a constituent of dietary fiber, linked to improve digestive

health, including increased gut motility and reduced risk of constipation.⁽²⁹⁾ Pomegranate peel's galacturonic acid exhibit immunomodulatory properties that may assist in infection prevention and bolster the immune system.⁽³¹⁾ Liquid holding capacity was evaluated and illustrated in Figure 2b. It was found that the extracted pectin was better at retaining water and dimethyl sulfoxide (DMSO) than acetic acid and acetone. A polymer with a higher ability to retain fluid can facilitate the production of low-calorie foods by enhancing volume without substantially increasing calorie content. Additionally, it directly impacts the food's texture and viscosity. As a result, the pomegranate peel pectin's capability to hold liquid is essential for food production from a technological and medicinal standpoint.^(29,32)





Figure 2. Physiochemical characteristics (a) and liquid holding capacity (b) of pectin derived from pomegranate (*Punica granatum* L.) peel

The FT-IR spectral analysis of the extracted pectin, as shown in **Figure 3**, revealed several characteristic absorption bands. A broad band between 3500-3300 cm^{-1} corresponds to non-associated hydroxyl (-OH) group. The absorbance near 2926 cm^{-1} is attributed to -CH, -CH₂, and -CH₃ stretching vibrations from galacturonic acid methyl esters. A distinct peak at 1740 cm^{-1} indicates C=O stretch, likely originating from esterified acetyl (COCH₃) groups. The peak observed at 1630 cm^{-1} is linked to the -OH tensile vibration band, while those at 1320 cm^{-1} and 1445 cm^{-1} suggest the presence of -CH₃ groups. Finally, the peak near 1026 cm^{-1} is associated with C-O stretching or bending vibrations. The structural characteristics observed in the FTIR spectrum, specifically methyl/acetyl

esterification and a resilient polysaccharide structure demonstrate the possible physiological and nutritional advantages of this pectin.

3.3 Physiochemical Properties of Pomegranate Juice

Pomegranate juice's pH was 3.2. The pH range of 3.1-3.3 is ideal for gel sets. Typically, gel formation is poor when the pH is above 3.5, while the gel is hard when the pH is below 3. According to the findings, fresh juice contains 85.3% moisture, 10.5% sugars (mostly fructose and glucose), 0.15g of total acidity (citric acid), 0.7mg of ascorbic acid, and 0.03g of ash per 100 mL. The chemical composition of pomegranate juice showed in Table 3 exhibit similarities to those reported in previous research by Sreekumar et al. (2014).⁽³³⁾

Table 3. Chemical composition of pomegranate juice

Parameters	Value*
pH	3.2±0.05
Moisture (%)	85.3±0.16
Total sugar (%)	10.5±0.02
Total acidity(g/100ml)	0.15±0.35
Ascorbic acid (mg/100ml)	0.9±0.03
Ash(g/100ml)	0.03±0.01

*Data expressed as mean ± standard deviation.

3.4 Physiochemical Analysis of Pomegranate Jelly

The pH, Acidity, and TSS of the five jelly samples found in this study are shown in Table 4. The table indicates a statistically significant difference in pH values among the samples, with readings from 2.81±0.01 to 3.02±0.01, which fall within the typical range for jellies. TSS values for pomegranate jelly in this study ranged from 0.62±0.01 to 0.67±0.02, which is roughly comparable to the values for dragon fruit jelly discovered by Islam et al. (2012).⁽³⁴⁾

The proximate makeup of the jelly samples is shown in Table 5. The table shows that, in contrast to the jellies created using extracted pectin, the control sample has a higher moisture content. The relatively low moisture content observed in samples b and e might be due to sugar, as sugar reduces the moisture content, which inhibits the growth of food-spoiling microbes. Samples PJ_c and PJ_d exhibit higher moisture content compared Samples PJ_b and PJ_e due to the presence of honey, which acts as a hydrophilic agent in low humid condition. The sample will therefore have a longer shelf life than the sample containing honey when stored under the same conditions due to its lower moisture content.⁽³⁵⁾ Fiber, ash, protein, and fat content do not have any significant differences from control PJ_a, indicating that extracted pectin possess a quality comparable to

commercial pectin, as pectin does not alter this composition. Samples PJ_b and PJ_e have higher carbohydrate content than samples PJ_c and PJ_d, as these have been prepared with sugar

Table 4. Physical parameters of jelly

Sample	pH	Acidity	TSS
PJ _a	2.93±0.02 ^b	0.64±0.01 ^b	0.62±0.01 ^c
PJ _b	3.02±0.01 ^a	1.03±0.01 ^a	0.67±0.02 ^a
PJ _c	3.00±0.02 ^a	1.00±0.02 ^a	0.66±0.06 ^a
PJ _d	2.81±0.01 ^c	0.60±0.01 ^{bc}	0.65±0.04 ^b
PJ _e	2.9±0.05 ^b	0.63±0.03 ^b	0.61±0.01 ^c
<i>p</i> -value	0.01	0.02	0.01

Values are presented as mean ± SD. Within the same column, means sharing identical superscripts do not differ significantly ($P < 0.05$). Where, PJ_a: Pomegranate jelly made with commercial pectin (control), PJ_b: 40% Sugar + 60% Pomegranate fruit juice + Extracted pectin, PJ_c: 40% Honey + 60% Pomegranate fruit juice + Extracted pectin, PJ_d: 50% Honey + 50% Pomegranate fruit juice + Extracted pectin, PJ_e: 50% Sugar + 50% Pomegranate fruit juice + Extracted pectin

3.5 Antioxidant Activity of Pomegranate Jelly

Antioxidant capacity of the prepared jelly is demonstrated in Figure 4, indicates that there were no significant differences between the control and other samples in terms of antioxidant capacity, which means that extracted pectin has increased antioxidant activity. The substantial antioxidant capacity of pomegranate jelly is due to its high concentration of polyphenols and flavonoids which provide protection against disorders associated to oxidative stress, including cardiovascular diseases, diabetes, and cancer.^(36,37) These findings highlight the functional dietary potential of pomegranate jellies; nevertheless, further real-time studies are required to confirm their bioavailability and long term health impacts.

Table 5. Chemical composition of pomegranate jelly (expressed as percentage, %)

Parameters	PJ _a	PJ _b	PJ _c	PJ _d	PJ _e
Moisture	36.76 ±0.01 ^a	30.70±0.01 ^c	33.82± 0.01 ^b	34.75±0.01 ^b	31.07±0.01 ^c
Crude Fibre	1.30±0.04 ^b	1.59±0.03 ^a	1.80±0.06 ^a	1.92±0.02 ^a	1.93± 0.01 ^a
Ash	0.81±0.03 ^a	1.61±0.02 ^a	1.30±0.01 ^a	1.37±0.03 ^a	0.71±0.06 ^a
Fat	0.92±0.02 ^a	1.20±0.09 ^a	1.58±0.06 ^a	1.79±0.02 ^a	1.35±0.03 ^a
Protein	1.58±0.03 ^a	1.50±0.05 ^a	1.98±0.02 ^a	1.87±0.01 ^a	1.40±0.06 ^a
Carbohydrate	58.63±0.06 ^b	64.32±0.04 ^a	59.52±0.03 ^b	58.3±0.01 ^b	63.54±0.02 ^a

Values are presented as mean ± SD. Within the same column, means sharing identical superscripts do not differ significantly ($p < 0.05$). Where, PJ_a: Pomegranate jelly made with commercial pectin (control), PJ_b: 40% Sugar + 60% Pomegranate fruit juice + Extracted pectin, PJ_c: 40% Honey + 60% Pomegranate fruit juice + Extracted pectin, PJ_d: 50% Honey + 50% Pomegranate fruit juice + Extracted pectin, PJ_e: 50% Sugar + 50% Pomegranate fruit juice + Extracted pectin




Figure 3. FT-IR spectrum of pectin obtained from dried pomegranate peel; Wavenumber (cm⁻¹) is ranged from 450 to 4000 cm⁻¹ (x axis); %T is the percentage of transmittance (y axis)




Figure 4. Antioxidant activity of pomegranate jelly

3.6 Microbial Analysis of Pomegranate Jelly

Up to 90 days of storage at 5°C, there was no growth of *E. coli* in the jelly sample. According to Pramanick et al. (2021), the complete absence of total coliform count, total fungal count, *E. coli*, *Streptococcus* spp., *Vibrio* spp. in jelly was an indication of better and safer quality jelly production.⁽³⁸⁾ Yeast and mold were not found in the jelly up to 30 days of storage time. After that

period, yeast and mold were grown gradually according to Table 6.

At 5°C storage conditions, the yeast and mold count of jelly was zero till 30 days. But it was increased slightly from 0.2×10³ cfu/g to 2×10³ cfu/g after 30 days. From previous study on wood apple jelly stored at ambient temperature conducted by Kumar and Deen (2017) showed 3×10³ cfu/g microbial counts.⁽³⁹⁾ Gaikwad (2016) also found similar results for sapota and beetroot

blended jelly, where the microbial count was 2×10^3 cfu/g.⁽³⁵⁾ In the present findings, the microbial count of sample P_{Jb} and P_{Je} had exceeded this limit means that after 60 days, the jelly could lose its acceptable organoleptic property. Despite having higher moisture

content than the control samples, honey-added jelly (Samples P_{Jc} and P_{Ja}) had a progressive decrease in microbial growth, suggesting that honey was inhibitory to the growth of microorganisms.

Table 6. Microbial count of jelly at 5°C up to 90 90-day storage period

Storage days	Yeast and mold (cfu/gm)				
	P _{Ja}	P _{Jb}	P _{Jc}	P _{Ja}	P _{Je}
0	Not found	Not found	Not found	Not found	Not found
30	Not found	Not found	Not found	Not found	Not found
60	$1 \pm 0.12 \times 10^{3a}$	$0.9 \pm 0.11 \times 10^{3c}$	$0.4 \pm 0.26 \times 10^{3b}$	$0.2 \pm 0.23 \times 10^{3b}$	$1 \pm 0.27 \times 10^{3a}$
90	$2 \pm 0.17 \times 10^{3a}$	$2 \pm 0.14 \times 10^{3a}$	$1 \pm 0.54 \times 10^{3c}$	$1 \pm 0.56 \times 10^{3c}$	$1.9 \pm 0.43 \times 10^{3b}$

P_{Ja}: Pomegranate jelly made with commercial pectin (control), P_{Jb}: 40% Sugar + 60% Pomegranate fruit juice + Extracted pectin, P_{Jc}: 40% Honey + 60% Pomegranate fruit juice + Extracted pectin, P_{Ja}: 50% Honey + 50% Pomegranate fruit juice + Extracted pectin, P_{Je}: 50% Sugar + 50% Pomegranate fruit juice + Extracted pectin

3.7 Sensory Evaluation

According to sensory analysis, it was interpreted that with an overall acceptance score of 5.5 ± 0.02 , the jelly made with additional sugar (sample P_{Je}) had the highest score of any formulation (Figure 5). This positive rating is

probably the result of its pleasing flavor, taste, color, texture, and appearance. Interestingly, sample P_{Je} performed similarly to the control (P_{Ja}). As illustrated in Figure 5, the honey-based samples P_{Jc} and P_{Ja}, on the other hand, obtained the lowest sensory scores, with ratings of 4.8 ± 0.01 and 4.5 ± 0.03 , respectively.




Figure 5. Radar chart of sensory properties. Values are means of 10 panelists. Where, P_{Ja}: Pomegranate jelly made with commercial pectin (control), P_{Jb}: 40% Sugar + 60% Pomegranate fruit juice + Extracted pectin, P_{Jc}: 40% Honey + 60% Pomegranate fruit juice + Extracted pectin, P_{Ja}: 50% Honey + 50% Pomegranate fruit juice + Extracted pectin, P_{Je}: 50% Sugar + 50% Pomegranate fruit juice + Extracted pectin

One possible explanation for the jelly’s deeper hue and slightly acidic flavor is the honey content, which was different from that of the control sample (P_{Ja}). Furthermore, the jelly’s flavor and taste were significantly shaped by the combination of pectin and citric acid. Increasing the volumes of gelatin and citric acid improve

the texture and consistency of the jelly, as reported by Basu and Shivhare (2010).⁽⁴⁰⁾

The study provides valuable insights into the physiochemical characteristics of pomegranate peel pectin to be used as a gelling agent in jelly formation. In comparison to the commercial pectin (control), the

extracted pectin formulated jelly showed enhanced nutritional composition, increased antioxidant activity, and favorable textural qualities. The findings suggest that pomegranate peel pectin can successfully substitute synthetic gelling agents while providing additional functional health benefits. However, certain limitations in this study must be acknowledged. The study was conducted at a laboratory setting, but specific clinical evidence is scarced, requiring further research to confirm the bioavailability, efficacy and potential health benefits of pomegranate peel pectin jelly in dietary applications. The sensory evaluation conducted in this study relied on a limited panel, which might not properly represent consumer diversity. Furthermore, for generating more robust proof for the incorporation of pomegranate peel pectin in functional foods and expand its utilization in food sector, future study should focus on investigating the bioavailability of functional compounds in human model.

4. CONCLUSION

Pomegranate peel pectin is a natural ingredient that has the potential to be a promising alternative to synthetic pectin in the production of jellies that provide additional nutritional benefits. From this study, it is evident that the pectin extracted from pomegranate peel exhibited jelly-forming properties similar to that of commercial pectin. It boosts the jelly's health promoting value by adding fiber and bioactive substances in addition to its gelling properties. The property of sugar-added pomegranate peel pectin jelly to remain consumable for up to three months at 5C confirmed its viability for both industrial and consumer use. Future studies should concentrate on sugar-reduces or sugar-free formulations, assessment of the bioavailability of bioactive compounds, and optimization for large-scale extraction. Determining its antioxidant, probiotic, and diseases preventing properties might enhance the role of pomegranate peel pectin as a sustainable by-product and a functional food component that promotes human health.

Ethical Approval

The sensory evaluation procedures of this study were conducted in accordance with the ethical guidelines of the Department of Applied Food Science and Nutrition, Faculty of Food Science and Technology, Chattogram Veterinary and Animal Sciences University (CVASU).

Ethical approval was granted under Ref. No. CVASU/Head/AFSN-01/2012/1276(1). All participants provided informed consent prior to participation and were informed of their right to withdraw at any time without consequence.

Acknowledgement

The authors sincerely express their acknowledgements to the Department of Applied Food Science and Nutrition, the Department of Applied Chemistry and Chemical Technology and the Department of Food Processing and Engineering, Chattogram Veterinary and Animal Sciences University (CVASU), for providing technical assistance and laboratory facilities throughout the research.

Competing Interests

All the authors declare that there are no conflicts of interest.

Funding Information

This work was funded by Ministry of Science and Technology, Bangladesh and Advanced Studies and Research (CASR), CVASU, Bangladesh. The authors are grateful for the financial support that made this research possible.

Underlying Data

Derived data supporting the findings of this study are available from the corresponding author on request.

REFERENCES

1. Baker MT, Lu P, Parrella JA, Leggette HR. Consumer acceptance toward functional foods: A scoping review. *International Journal of Environmental Research and Public Health*. 2022;19(3):1217. <https://doi.org/10.3390/ijerph19031217>
2. Mo Y, Ma J, Gao W, Zhang L, Li J, Li J, Zang J. Pomegranate peel as a source of bioactive compounds: A mini review on their physiological functions. *Frontiers in Nutrition*. 2022;9:887113. <https://doi.org/10.3389/fnut.2022.887113>
3. Kshirsagar KR, Pathak SS, Patil SM. Pomegranate (*Punica granatum* L.): A fruitful fountain of remedial potential. *Cureus*. 2023;15(9):e45677. <http://dx.doi.org/10.7759/cureus.45677>
4. Mohan M, Mohanavarshaa CA, Priya D. Review of pharmacological and medicinal uses of *punica granatum*. *Cureus*. 2024;16(10). <https://doi.org/10.7759/cureus.71510>
5. Dogara AM, Hama HA, Ozdemir D. Update on the potential of *Punica granatum* L. traditional uses and

- pharmacological uses: A review. *Advances in Pharmacological and Pharmaceutical Sciences*. 2024;2024(1):6523809. <https://doi.org/10.1155/adpp/6523809>
6. Sarkar A, Haque MA, Alam M. Unlocking the potential of pomegranate peels as a valuable source of bioactive compounds through effective drying strategies. *Food Chemistry Advances*. 2024 1;4:100622. <https://doi.org/10.1016/j.focha.2024.100622>
 7. Abid M, Cheikhrouhou S, Renard CM, Bureau S, Cuvelier G, Attia H, Ayadi MA. Characterization of pectins extracted from pomegranate peel and their gelling properties. *Food Chemistry*. 2017;215:318-325. <https://doi.org/10.1016/j.foodchem.2016.07.181>
 8. Dambuza A, Rungqu P, Oyedeji AO, Miya G, Oriola AO, Hosu YS, Oyedeji OO. Therapeutic potential of pectin and its derivatives in chronic diseases. *Molecules*. 2024;29(4):896. <https://doi.org/10.3390/molecules29040896>
 9. Yi L, Cheng L, Yang Q, Shi K, Han F, Luo W, Duan S. Source, extraction, properties, and multifunctional applications of pectin: A short review. *Polymers*. 2024;16(20):2883. <https://doi.org/10.3390/polym16202883>
 10. Soomro MA, Khan S, Majid A, Bhatti S, Perveen S, Phull AR. Pectin as a biofunctional food: comprehensive overview of its therapeutic effects and antidiabetic-associated mechanisms. *Discover Applied Sciences*. 2024;6(6):298. <http://dx.doi.org/10.1007/s42452-024-05968-1>
 11. Gonzales E, Bustamante A, Garcia-Diaz D, Sanhueza L, Orellana JF, Fredes C, Jimenez P, Chavez V, Echeverría F. Assessing quality and polyphenol in vitro bioaccessibility in healthy jelly gummies with microencapsulated and non-encapsulated pomegranate peel extract. *Food Chemistry*. 2025;470:142611. <https://doi.org/10.1016/j.foodchem.2024.142611>
 12. Horison R, Surini S. Utilization of glucomannan-xanthan gum-carrageenan based co-processed excipients in the development of nutraceutical pomegranate (*Punica granatum L.*) peel extract jelly. *Journal of Pharmacy & Pharmacognosy Research*. 2025;13(3):774–786. http://dx.doi.org/10.56499/jppres24.2060_13.3.774
 13. Mahmoud MH, Azab DE. Upcycling mango and pomegranate peel as pectin source: characterization, fat substitute and retarded lipid oxidation in caramel candy. *Discover Food*. 2025;5(1):37. <https://doi.org/10.1007/s44187-025-00295-9>
 14. Firat E, Koca N, Kaymak-Ertekin F. Extraction of pectin from watermelon and pomegranate peels with different methods and its application in ice cream as an emulsifier. *Journal of Food Science*. 2023;88(11):4353-4374. <https://doi.org/10.1111/1750-3841.16752>
 15. Diaz-Herrera R, Alvarez-Pérez OB, Ventura-Sobrevilla J, Ascacio-Valdés A, Aguilar-Gonzalez MA, Buenrostro-Figueroa J, Aguilar CN. Pomegranate peel polyphenols as an antioxidant additive for the development and characterization of a new active pectin edible film. *eFood*. 2023;4(6):e115. <https://doi.org/10.1002/efd2.115>
 16. Sathish S, Gowthaman KA, Augustus H. Utilization of *Punica granatum* peels for the extraction of pectin. *Research Journal of Pharmacy and Technology*. 2018;11(2):613-616. <http://dx.doi.org/10.5958/0974-360X.2018.00113.0>
 17. Aina VO, Barau MM, Mamman OA, Zakari A, Haruna H, Umar MH, Abba YB. Extraction and characterization of pectin from peels of lemon (*Citrus limon*), grapefruit (*Citrus paradisi*), and sweet orange (*Citrus sinensis*). *British Journal of Pharmacology and Toxicology*. 2012;3(6):259-262.
 18. AOAC. Official Methods of Analysis of the Association of Official Analytical Chemists; 2000. Gaithersburg (MD): AOAC International; 2000.
 19. FAO JECFA. Compendium of food additive specifications: 71st meeting. Rome: Food and Agriculture Organization of the United Nations; 2009.
 20. Zhang L, Zhu M, Shi T, Guo C, Huang Y, Chen Y, Xie M. Recovery of dietary fiber and polyphenol from grape juice pomace and evaluation of their functional properties and polyphenol compositions. *Food & Function*. 2017;8(1):341-351. <https://doi.org/10.1039/c6fo01423b>
 21. Wathoni N, Shan CY, Shan WY, Rostinawati T, Indradi RB, Pratiwi R, Muchtaridi M. Characterization and antioxidant activity of pectin from Indonesian mangosteen (*Garcinia mangostana L.*) rind. *Heliyon*. 2019;5(8). <https://doi.org/10.1016/j.heliyon.2019.e02299>
 22. Azlim Almey AA, Ahmed Jalal Khan C, Syed Zahir I, Mustapha Suleiman K, Aisyah MR, Kamarul Rahim K. Total phenolic content and primary antioxidant activity of methanolic and ethanolic extracts of aromatic plants' leaves. *International Food Research Journal*. 2010;17(4):1077-1084.
 23. Aryal S. Sabouraud Dextrose Agar (SDA)–Composition, Principle, Uses, Preparation and Colony Morphology. 2015. Available from: <https://microbiologyinfo.com/sabouraud-dextrose-agar-sda-composition-principle-uses-preparation-and-colony-morphology/> (Accessed on 14 Jul 2025)
 24. Peterkin PI. Compendium of methods for the microbiological examination of foods (3rd edn). *Trends in Food Science & Technology*. 1993;4(6):199. [http://dx.doi.org/10.1016/0924-2244\(93\)90131-s](http://dx.doi.org/10.1016/0924-2244(93)90131-s)
 25. US Food and Drug Administration (USFDA), Center for Food Safety and Applied Nutrition (CFSAN). Rapid methods for detecting foodborne pathogens. *Bacteriological Analytical Manual Online*. 2001. Available from: <http://www.fda.gov/downloads/Food/FoodScienceResearch/UCM244777.pdf> (Accessed on 14 Jul 2025)
 26. Yuliarti O, Goh KK, Matia-Merino L, Mawson J, Brennan C. Extraction and characterisation of pomace pectin from gold kiwifruit (*Actinidia chinensis*). *Food Chemistry*. 2015;187:290–296. <https://doi.org/10.1016/j.foodchem.2015.03.148>
 27. Panchami PS, Gunasekaran S. Extraction and characterization of pectin from fruit waste. *International Journal of Current Microbiology and Applied Sciences*. 2017;6(8):943-948. <http://dx.doi.org/10.20546/ijcmas.2017.608.116>
 28. Mollea C, Chiampo F, Conti R. Extraction and

- characterization of pectins from cocoa husks: A preliminary study. *Food Chemistry*. 2008;107(3):1353–1356. <http://dx.doi.org/10.1016/j.foodchem.2007.09.006>
29. Dixit SS, Muruganandam L, Moorthy IG. Pectin from fruit peel: A comprehensive review on various extraction approaches and their potential applications in pharmaceutical and food industries. *Carbohydrate Polymer Technologies and Applications*. 2025;9:100708. <https://doi.org/10.1016/j.carpta.2025.100708>
30. Georgiev Y, Ognyanov M, Yanakieva I, Kussovski V. Isolation, characterization and modification of citrus pectins. *Journal of BioScience and Biotechnology*. 2012;1(3):223-233.
31. Kandylis P, Kokkinomagoulos E. Food applications and potential health benefits of pomegranate and its derivatives. *Foods*. 2020;9(2):122. <https://doi.org/10.3390/foods9020122>
32. Rubio-Senent F, Rodríguez-Gutiérrez G, Lama-Muñoz A, Fernández-Bolaño J. Pectin extracted from thermally treated olive oil by-products: Characterization, physico-chemical properties, in vitro bile acid and glucose binding. *Food Hydrocoll*. 2015;43:311–321. <http://dx.doi.org/10.1016/j.foodhyd.2014.06.001>
33. Sreekumar S, Sithul H, Muraleedharan P, Azeez JM, Sreeharshan S. Pomegranate fruit as a rich source of biologically active compounds. *BioMed Research International*. 2014;2014(1):686921. <https://doi.org/10.1155/2014/686921>
34. Islam M, Khan M, Hoque M, Rahman M. Studies on the Processing and Preservation of Dragon Fruit (*Hylocereus undatus*) Jelly. *The Agriculturists*. 2012;10(2):29–35. <http://dx.doi.org/10.3329/agric.v10i2.13139>
35. Gaikwad SU. Studies on preparation of sapota (*Manilkara achras*)–beetroot (*Beta vulgaris*) blended jelly [Master's thesis]. Maharashtra, India: PG Institute of Post-Harvest Management; 2016.
36. Saparbekova AA, Kantureyeva GO, Kudasova DE, Konarbayeva ZK, Latif AS. Potential of phenolic compounds from pomegranate (*Punica granatum L.*) by-product with significant antioxidant and therapeutic effects: A narrative review. *Saudi journal of biological sciences*. 2023 Feb 1;30(2):103553.
37. Parisi V, Santoro V, Donadio G, Bellone ML, Diretto G, Sandri C, Mensitieri F, De Tommasi N, Dal Piaz F, Braca A. Comparative chemical analysis of eight *Punica granatum L.* Peel cultivars and their antioxidant and anti-inflammatory activities. *Antioxidants*. 2022;11(11):2262. <https://doi.org/10.3390/antiox11112262>
38. Pramanick P, Zaman S, Mitra A. Mangrove fruit-based jelly as roadmap to alternative livelihood for island dwellers of Indian Sundarbans. In: Mitra A, Calma MM, Chakrabarty SP. *Natural Resources And Their Ecosystem Services (Vol. II)*. Bangalore, India: HSRA Publication 2021. p. 121–126.
39. Kumar A, Deen B. Studies on preparation and storage of jelly from wood apple (*Limonia acidissima L.*) fruits. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(6):224–229.
40. Islam M, Khan M, Hoque M, Rahman M. Studies on the Processing and Preservation of Dragon Fruit (*Hylocereus undatus*) Jelly. *The Agriculturists*. 2012;10(2):29–35. <http://dx.doi.org/10.3329/agric.v10i2.13139>